



Poisson Lie Sigma Models

Boris Arm

► To cite this version:

| Boris Arm. Poisson Lie Sigma Models. 2012. hal-00809008

HAL Id: hal-00809008

<https://hal.science/hal-00809008>

Preprint submitted on 8 Apr 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Some Poisson-Lie sigma models

ARM Boris

arm.boris@gmail.com

Abstract

We calculate the Poisson-Lie sigma model for every 4-dimensional Manin triples (function of its structure constant) and we give the 6-dimensional models for the Manin triples

$$\begin{aligned} &(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}), \mathfrak{sl}(2, \mathbb{C})^*), \\ &(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}), \\ &(\mathfrak{sl}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C})) \text{ and} \\ &(\mathfrak{sl}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C})) \end{aligned}$$

1 Introduction

A Manin triples $(\mathfrak{D}, \mathfrak{g}, \tilde{\mathfrak{g}})$ is a bialgebra $(\mathfrak{g}, \tilde{\mathfrak{g}}$ which don't intersect each others and a direct sum of this bialgebra $\mathfrak{D} = \mathfrak{g} \oplus \tilde{\mathfrak{g}}$). If the corresponding Lie groups have a Poisson structure, they are called Poisson-Lie groups. A Poisson-Lie sigma models is an action (3.13) calculated by a Poisson vector field matrix. [3] have deduced the extremal field which minimize the action of this models, which gives the motion equation (3.19). We calculate here the action and the equations of motion for some 6-dimensionals Manin triples and we give a general formula for each 4-dimensional Manin triples. The 6-dimensional Manin triples are $(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}), \mathfrak{sl}(2, \mathbb{C})^*), (\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}), \mathfrak{sl}(2, \mathbb{C})^*), (\mathfrak{sl}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C}))$ and $(\mathfrak{sl}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C}))$.

2 Some Manin triples

The Drinfeld double D is defined as a Lie group such that its Lie algebra \mathfrak{D} equipped by a symmetric ad-invariant nondegenerate bilinear form $\langle ., . \rangle$ can be decomposed into a pair of maximally isotropic subalgebras $\mathfrak{g}, \tilde{\mathfrak{g}}$ such that \mathfrak{D} as a vector space is the direct sum of \mathfrak{g} and $\tilde{\mathfrak{g}}$. Any such decomposition written as an ordered set $(\mathfrak{D}, \mathfrak{g}, \tilde{\mathfrak{g}})$ is called a Manin triples $(\mathfrak{D}, \mathfrak{g}, \tilde{\mathfrak{g}}), (\mathfrak{D}, \tilde{\mathfrak{g}}, \mathfrak{g})$.

One can see that the dimensions of the subalgebras are equal and that bases $\{T_i\}, \{\tilde{T}^i\}$ in the subalgebras can be chosen so that

$$\langle T_i, T_j \rangle = 0, \quad \langle T_i, \tilde{T}^j \rangle = \langle \tilde{T}^j, T_i \rangle = \delta_i^j, \quad \langle \tilde{T}^i, \tilde{T}^j \rangle = 0 \quad (2.1)$$

This canonical form of the bracket is invariant with respect to the transformations

$$T'_i = T_k A_i^k, \quad \tilde{T}'^j = (A^{-1})^j_k \tilde{T}^k \quad (2.2)$$

Due to the ad-invariance of $\langle ., . \rangle$ the algebraic structure of \mathfrak{D} is

$$\begin{aligned} [T_i, T_j] &= c_{ij}^k T_k, & [\tilde{T}^i, \tilde{T}^j] &= f^{ij}_k \tilde{T}^k \\ [T_i, \tilde{T}^j] &= f^{jk}_i T_k - c_{ik}^j \tilde{T}^k \end{aligned}$$

There are just four types of nonisomorphic four-dimensional Manin triples.

Abelian Manin triples :

$$[T_i, T_j] = 0, \quad [\tilde{T}^i, \tilde{T}^j] = 0, \quad [T_i, \tilde{T}^j] = 0, \quad i, j = 1, 2 \quad (2.3)$$

Semi-Abelian Manin triples (only non trivial brackets are displayed) :

$$[\tilde{T}^1, \tilde{T}^2] = \tilde{T}^2, \quad [T_2, \tilde{T}^1] = T_2, \quad [T_2, \tilde{T}^2] = -T_1 \quad (2.4)$$

Type A non-Abelian Manin triples ($\beta \neq 0$) :

$$\begin{aligned} [T_1, T_2] &= T_2, & [\tilde{T}^1, \tilde{T}^2] &= \beta \tilde{T}^2 \\ [T_1, \tilde{T}^2] &= -\tilde{T}^2, & [T_2, \tilde{T}^1] &= \beta T_2, & [T_2, \tilde{T}^2] &= -\beta T_1 + \tilde{T}^1 \end{aligned}$$

Type B non-Abelian Manin triples :

$$\begin{aligned} [T_1, T_2] &= T_2, & [\tilde{T}^1, \tilde{T}^2] &= \tilde{T}^1 \\ [T_1, \tilde{T}^1] &= T_2, & [T_1, \tilde{T}^2] &= -T_1 - \tilde{T}^2, & [T_2, \tilde{T}^2] &= \tilde{T}^1 \end{aligned}$$

Now we focus some six dimensional Manin triples. We recall that the commutation relations of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ of the Lie group $SL(2, \mathbb{C})$:

$$[T_1, T_2] = 2T_2, \quad [T_1, T_3] = -2T_3, \quad [T_2, T_3] = T_1 \quad (2.5)$$

The dual Lie algebra $\mathfrak{sl}(2, \mathbb{C})^*$ of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ has the commutation relations :

$$[\tilde{T}^1, \tilde{T}^2] = \frac{1}{4} \tilde{T}^2, \quad [\tilde{T}^1, \tilde{T}^3] = \frac{1}{4} \tilde{T}^3, \quad [\tilde{T}^2, \tilde{T}^3] = 0 \quad (2.6)$$

There is a scalar product on $(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*)$ such that (see [2]) :

$$(T_i, \tilde{T}^j) = \delta_i^j \quad (2.7)$$

Finally, we have that $(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}), \mathfrak{sl}(2, \mathbb{C})^*)$ with this scalar product is a Manin triple. We note that $(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}))$ with this scalar product is also a Manin triples.

The Iwasawa decomposition allows us to decompose :

$$\mathfrak{sl}(2, \mathbb{C}) = \mathfrak{su}(2, \mathbb{C}) \oplus \mathfrak{sb}(2, \mathbb{C}) \quad (2.8)$$

where $\mathfrak{su}(2, \mathbb{C})$ is the Lie algebra of the Lie group $SU(2)$ with commutation relations :

$$[T_1, T_2] = T_3, \quad [T_2, T_3] = T_1, \quad [T_3, T_1] = T_2 \quad (2.9)$$

$\mathfrak{sb}(2, \mathbb{C})$ is the Lie algebra of the Borel subgroup $SB(2, \mathbb{C})$ with commutation relations :

$$[\tilde{T}^1, \tilde{T}^2] = \tilde{T}^2, \quad [\tilde{T}^1, \tilde{T}^3] = \tilde{T}^3, \quad [\tilde{T}^2, \tilde{T}^3] = 0 \quad (2.10)$$

Here we can see in comparing (2.10) and (2.6) that $\mathfrak{sb}(2, \mathbb{C}) \simeq \mathfrak{sl}(2, \mathbb{C})^*$.

The Iwasawa decomposition (2.8) allows us to identify $\mathfrak{sb}(2, \mathbb{C}) \simeq \mathfrak{su}(2, \mathbb{C})^*$. We define a scalar product on $(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*)$ such that $(x, y) = \text{Im}(\text{Tr}(x|y))$. With this scalar product we have (see [2]) :

$$(T_i, \tilde{T}^j) = \delta_i^j \quad (2.11)$$

Finally we have that $(\mathfrak{sl}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C}))$ with this scalar product is a Manin triple. We note that $(\mathfrak{sl}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C}))$ with this scalar product is also a Manin triples.

3 Poisson-Lie sigma models

Given a Lie group M and a Poisson structure on it. We define the action of this model (see [1]) as :

$$S_1 = \int_{\Sigma} (< dg g^{-1}, A > - \frac{1}{2} < A, (r - \text{Ad}_g r \text{Ad}_g) A >) \quad (3.12)$$

where $g \in G$, $A = A_{\alpha}^i d\xi^{\alpha} X_i$ and $r \in \mathfrak{g} \otimes \mathfrak{g}$ is a classical r matrix with \mathfrak{g} as the Lie algebra of G and $\{X_i\}$ as a basis of \mathfrak{g} . Note that the above action can be applied for simple or nonsemisimple Lie group G with ad-invariant symmetric bilinear nondegenerate form $< X_i, X_j > = G_{ij}$ on the Lie algebra \mathfrak{g} . When the metric G_{ij} of Lie algebra is denegerate then the above action is not good. Here we use the following action instead of the above one :

$$S_2 = \int_{\Sigma} (dX_i \wedge A_i - \frac{1}{2} \mathcal{P}^{ij} A_i \wedge A_j) \quad (3.13)$$

where x are Lie group parameters with parametrization (e.g.)

$$\forall g \in G, g = e^{X_1 T_1} e^{X_2 T_2} \dots \quad (3.14)$$

where P^{ij} is the Poisson structure on the Lie group which for coboundary Poisson Lie groups it is obtained from

$$(\mathcal{P}(g))_\chi = b(g)a(g)^{-1} \quad (3.15)$$

We can obtain $a(g)^{-1}$ and $b(g)$ in computing :

$$(Ad_{g^{-1}})_\chi = \begin{pmatrix} a(g)^T & b(g)^T \\ 0 & d(g)^T \end{pmatrix} \quad (3.16)$$

$$(Ad_g)_\chi = \begin{pmatrix} a(g)^{-T} & -a(g)^{-T}b(g)^Td(g)^{-T} \\ 0 & d(g)^{-T} \end{pmatrix} \quad (3.17)$$

where T denotes the transpose matrix.

The extremal fields (X, A) which minimize the action (3.13) have to satisfy the equation written locally (see [3]) as :

$$dX_i + \mathcal{P}^{ij}(X)A_j = 0 \quad (3.18)$$

$$dA_k + \frac{1}{2}\mathcal{P}^{ij}_{,k}(X)A_i \wedge A_j = 0 \quad (3.19)$$

where $\mathcal{P}^{ij}_{,k} = \partial_k \mathcal{P}^{ij}|_{X_k=0}$.

4 Poisson-Lie sigma model of any 4-dimensional Manin triple

We first calculate the matrix of the adjoint actions function of structure constant :

$$\begin{aligned} ad_{T_1} &= \begin{pmatrix} 0 & c_{12}^1 & 0 & -f^{12}_1 \\ 0 & c_{12}^2 & f^{12}_1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -c_{12}^1 & -c_{12}^2 \end{pmatrix} \\ ad_{T_2} &= \begin{pmatrix} -c_{12}^1 & 0 & 0 & -f^{12}_2 \\ -c_{12}^2 & 0 & f^{12}_2 & 0 \\ 0 & 0 & c_{12}^1 & c_{12}^2 \\ 0 & 0 & 0 & 0 \end{pmatrix} \end{aligned}$$

To obtain the matrix \mathcal{P} , we calculate the adjoint action matrix of a general element $g = \prod_{i=1}^2 e^{\alpha_i T_i}$ by the formula :

$$(Ad_{\prod_{i=1}^2 e^{X_i T_i}})_\chi = \prod_{i=1}^2 e^{X_i (ad_{T_i})_\chi} \quad (4.20)$$

Similarly, we have :

$$(Ad_{(\prod_{i=1}^2 e^{X_i T_i})^{-1}})_\chi = \prod_{i=1}^2 e^{-X_{3-i} (ad_{T_{3-i}})_\chi} \quad (4.21)$$

We can deduce the matrix \mathcal{P}^{ij} :

$$\mathcal{P}^{ij} = \begin{pmatrix} 0 & -\mathcal{P}^{21} \\ \mathcal{P}^{21} & 0 \end{pmatrix}$$

where

$$\mathcal{P}^{21} = \frac{c_{12}^{-1}(-1 + e^{c_{12}^2 X_1})f_{12}^{12} + c_{12}^2 e^{c_{12}^2 X_1 - c_{12}^{-1} X_2}(-1 + e^{c_{12}^{-1} X_2})f_{12}^{12}}{c_{12}^2 c_{12}^{-1}} \quad (4.22)$$

Now, we can calculate the action (3.13) of the model

$$S_2 = \int_{\Sigma} \sum_{i=1}^2 dX_i \wedge A_i - \mathcal{P}^{21} A_2 \wedge A_1 \quad (4.23)$$

and the equations of motion (3.19) :

$$\begin{aligned} dX_1 - \mathcal{P}^{21} A_2 &= 0 \\ dX_2 + \mathcal{P}^{21} A_1 &= 0 \\ dA_1 - \frac{c_{12}^{-1} c_{12}^2 f_{12}^{12} + c_{12}^2 c_{12}^2 (-e^{-c_{12}^{-1} X_2} + 1) f_{12}^{12}}{c_{12}^2 c_{12}^{-1}} A_1 \wedge A_2 &= 0 \\ dA_2 + \frac{c_{12}^2 e^{c_{12}^2 X_1} c_{12}^{-1} f_{12}^{12}}{c_{12}^2 c_{12}^{-1}} A_2 \wedge A_1 &= 0 \end{aligned} \quad (4.24)$$

5 Poisson-Lie sigma model of $(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}), \mathfrak{sl}(2, \mathbb{C})^*)$

We first calculate the matrix of the adjoint actions :

$$\begin{aligned} ad_{T_1} &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix} \\ ad_{T_2} &= \frac{1}{4} \begin{pmatrix} 0 & 0 & 4 & 0 & -1 & 0 \\ -8 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 8 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -4 & 0 & 0 \end{pmatrix} \\ ad_{T_3} &= \frac{1}{4} \begin{pmatrix} 0 & -4 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 8 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -8 \\ 0 & 0 & 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{aligned}$$

To obtain the matrix \mathcal{P} , we calculate the adjoint action matrix of a general element $g = \prod_{i=1}^3 e^{\alpha_i T_i}$ by the formula :

$$(Ad_{\prod_{i=1}^3 e^{X_i T_i}})_\chi = \prod_{i=1}^3 e^{X_i (ad_{T_i})_\chi} \quad (5.25)$$

Similarly, we have :

$$(Ad_{(\prod_{i=1}^3 e^{X_i T_i})^{-1}})_\chi = \prod_{i=1}^3 e^{-X_i (ad_{T_i})_\chi} \quad (5.26)$$

We can deduce the matrix \mathcal{P}^{ij} :

$$\mathcal{P}^{ij} = \begin{pmatrix} 0 & -\frac{X_2}{4}(1 + X_2 X_3)e^{2X_1} & -\frac{X_3}{4}e^{-2X_1} \\ \frac{X_2}{4}(1 + X_2 X_3)e^{2X_1} & 0 & \frac{X_2 X_3}{2} \\ \frac{X_3}{4}e^{-2X_1} & -\frac{X_2 X_3}{2} & 0 \end{pmatrix}$$

Now, we can calculate the action (3.13) of the model

$$S_2 = \int_{\Sigma} \sum_{i=1}^3 dX_i \wedge A_i + \left(\frac{X_2}{4}(1 + X_2 X_3)e^{2X_1} \right) A_1 \wedge A_2 + \frac{X_3}{4}e^{-2X_1} A_1 \wedge A_3 - \frac{X_2 X_3}{2} A_2 \wedge A_3 \quad (5.27)$$

and the equations of motion (3.19) :

$$\begin{aligned} dX_1 - \left(\frac{X_2}{4}(1 + X_2 X_3)e^{2X_1} \right) A_2 - \frac{X_3}{4}e^{-2X_1} A_3 &= 0 \\ dX_2 + \left(\frac{X_2}{4}(1 + X_2 X_3)e^{2X_1} \right) A_1 + \frac{X_2 X_3}{2} A_3 &= 0 \\ dX_3 + \frac{X_3}{4}e^{-2X_1} A_1 - \frac{X_2 X_3}{2} A_2 &= 0 \\ dA_1 - \frac{X_2}{2}(1 + X_2 X_3) A_1 \wedge A_2 + \frac{X_3}{2} A_1 \wedge A_2 &= 0 \\ dA_2 - \frac{e^{2X_1}}{4} A_1 \wedge A_2 + \frac{X_3}{2} A_2 \wedge A_3 &= 0 \\ dA_3 - \frac{X_2^2}{4}e^{2X_1} A_1 \wedge A_2 - \frac{e^{-2X_1}}{4} A_1 \wedge A_3 + X_2 A_2 \wedge A_3 &= 0 \end{aligned}$$

6 Poisson-Lie sigma model of $(\mathfrak{sl}(2, \mathbb{C}) \oplus \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C})^*, \mathfrak{sl}(2, \mathbb{C}))$

Now to obtain this Poisson Lie sigma model, we have to change $T_i \rightarrow \tilde{T}^i$ and $\tilde{T}^i \rightarrow T_i$ of the previous model. And we can calculate the matrix of the adjoint actions as we do previously. With this we can deduce the matrix \mathcal{P}^{ij} for this model :

$$\mathcal{P}^{ij} = \begin{pmatrix} 0 & -2e^{\frac{X_1}{4}} X_2 & 2e^{\frac{X_1}{4}} X_3 \\ 2e^{\frac{X_1}{4}} X_2 & 0 & 2 - \frac{1}{2}e^{\frac{X_1}{2}}(4 + X_2 X_3) \\ -2e^{\frac{X_1}{4}} X_3 & -2 + \frac{1}{2}e^{\frac{X_1}{2}}(4 + X_2 X_3) & 0 \end{pmatrix}$$

Now, we can calculate the action (3.13) of the model

$$S_2 = \int_{\Sigma} \sum_{i=1}^3 dX_i \wedge A_i + 2e^{\frac{X_1}{4}} X_2 A_1 \wedge A_2 - 2e^{\frac{X_1}{4}} X_3 A_1 \wedge A_3 + (-2 + \frac{1}{2}e^{\frac{X_1}{2}}(4 + X_2 X_3)) A_2 \wedge A_3 \quad (6.28)$$

and the equations of motion (3.19) :

$$\begin{aligned} dX_1 - 2e^{\frac{X_1}{4}} X_2 A_2 + 2e^{\frac{X_1}{4}} X_3 A_3 &= 0 \\ dX_2 + 2e^{\frac{X_1}{4}} X_2 A_1 + (2 - \frac{1}{2}e^{\frac{X_1}{2}}(4 + X_2 X_3)) A_3 &= 0 \\ dX_3 - 2e^{\frac{X_1}{4}} X_3 A_1 - (2 - \frac{1}{2}e^{\frac{X_1}{2}}(4 + X_2 X_3)) A_2 &= 0 \\ dA_1 - \frac{X_2}{2} A_1 \wedge A_2 + \frac{X_3}{4} A_1 \wedge A_3 - \frac{1}{4}(4 + X_2 X_3) A_2 \wedge A_3 &= 0 \\ dA_2 - 2e^{\frac{X_1}{4}} A_1 \wedge A_2 - \frac{1}{2}e^{\frac{X_1}{4}} A_2 \wedge A_3 &= 0 \\ dA_3 + 2e^{\frac{X_1}{4}} A_1 \wedge A_3 - \frac{e^{\frac{X_1}{2}} X_2}{2} A_2 \wedge A_3 &= 0 \end{aligned}$$

7 Poisson-Lie sigma model of $(\mathfrak{sl}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C}))$

We can calculate the matrix of the adjoint actions as we do previously. With this we can deduce the matrix \mathcal{P}^{ij} for this model :

$$\mathcal{P}^{ij} = \begin{pmatrix} 0 & -\cos X_1 \cos X_3 \sin X_2 + \sin X_1 \sin X_3 & -\cos X_3 \sin X_1 \sin X_2 - \cos X_1 \sin X_3 \\ \cos X_1 \cos X_3 \sin X_2 - \sin X_1 \sin X_3 & 0 & -1 + \cos X_2 \cos X_3 \\ \cos X_3 \sin X_1 \sin X_3 + \cos X_1 \sin X_3 & 1 - \cos X_2 \cos X_3 & 0 \end{pmatrix}$$

Now, we can calculate the action (3.13) of the model

$$\begin{aligned} S_2 = \int_{\Sigma} \sum_{i=1}^3 dX_i \wedge A_i - (-\cos X_1 \cos X_3 \sin X_2 + \sin X_1 \sin X_3) A_1 \wedge A_2 \\ - (-\cos X_3 \sin X_1 \sin X_2 - \cos X_1 \sin X_3) A_1 \wedge A_3 - (-1 + \cos X_2 \cos X_3) A_2 \wedge A_3 \end{aligned} \quad (7.29)$$

and the equations of motion (3.19) :

$$\begin{aligned} dX_1 + (-\cos X_1 \cos X_3 \sin X_2 + \sin X_1 \sin X_3) A_2 + (-\cos X_3 \sin X_1 \sin X_2 - \cos X_1 \sin X_3) A_3 &= 0 \\ dX_2 + (\cos X_1 \cos X_3 \sin X_2 - \sin X_1 \sin X_3) A_1 + (-1 + \cos X_2 \cos X_3) A_3 &= 0 \\ dX_3 + (\cos X_3 \sin X_1 \sin X_3 + \cos X_1 \sin X_3) A_1 (1 - \cos X_2 \cos X_3) A_2 &= 0 \\ dA_1 + \sin X_3 A_1 \wedge A_2 - \cos X_3 \sin X_2 A_1 \wedge A_3 &= 0 \\ dA_2 - \cos X_1 \cos X_3 A_1 \wedge A_2 - \cos X_3 \sin X_1 A_1 \wedge A_3 &= 0 \\ dA_3 + \sin X_1 A_1 \wedge A_2 - \cos X_1 A_1 \wedge A_3 &= 0 \end{aligned}$$

8 Poisson-Lie sigma model of $(\mathfrak{sl}(2, \mathbb{C}), \mathfrak{sb}(2, \mathbb{C}), \mathfrak{su}(2, \mathbb{C}))$

We can calculate the matrix of the adjoint actions as we do previously. With this we can deduce the matrix \mathcal{P}^{ij} for this model :

$$\mathcal{P}^{ij} = \begin{pmatrix} 0 & -e^{X_1} X_3 & -e^{X_1} X_2 \\ e^{X_1} X_3 & 0 & \frac{1}{2}(1 - e^{2X_1}(1 + X_2^2 + X_3^2)) \\ e^{X_1} X_2 & \frac{1}{2}(-1 + e^{2X_1}(1 + X_2^2 + X_3^2)) & 0 \end{pmatrix}$$

Now, we can calculate the action (3.13) of the model

$$S_2 = \int_{\Sigma} \sum_{i=1}^3 dX_i \wedge A_i + e^{X_1} X_3 A_1 \wedge A_2 + e^{X_1} X_2 A_1 \wedge A_3 - \frac{1}{2}(1 - e^{2X_1}(1 + 2X_2^2 + 2X_3^2)) A_2 \wedge A_3 \quad (8.30)$$

and the equations of motion (3.19) :

$$\begin{aligned} dX_1 - e^{X_1} X_3 A_2 - e^{X_1} X_2 A_3 &= 0 \\ dX_2 + e^{X_1} X_3 A_1 + \frac{1}{2}(1 - e^{2X_1}(1 + X_2^2 + X_3^2)) A_3 &= 0 \\ dX_3 + e^{X_1} X_2 A_1 - \frac{1}{2}(1 - e^{2X_1}(1 + X_2^2 + X_3^2)) A_2 &= 0 \\ dA_1 - X_3 A_1 \wedge A_2 - X_2 A_1 \wedge A_3 - (1 + X_2^2 + X_3^2) A_2 \wedge A_3 &= 0 \\ dA_2 - e^{X_1} A_1 \wedge A_3 &= 0 \\ dA_3 - e^{X_1} A_1 \wedge A_2 &= 0 \end{aligned}$$

9 Discussion

We gives here the Poisson-Lie sigma models of some Manin triples. Concerning the general formula (9.32), we have to say that this is no problem when c_{12}^2 and c_{12}^2 is zero because

$$\mathcal{P}^{21} = \frac{(-1 + e^{c_{12}^2 X_1}) f_1^{12}}{c_{12}^2} + \frac{e^{c_{12}^2 X_1 - c_{12}^1 X_2} (-1 + e^{c_{12}^1 X_2}) f_2^{12}}{c_{12}^2} \quad (9.31)$$

which can be approximate by

$$\mathcal{P}^{21} = (X_1 + \frac{c_{12}^2}{2} X_1^2 + \dots) f_1^{12} + e^{c_{12}^2 X_1 - c_{12}^1 X_2} (X_2 + \frac{c_{12}^1}{2} X_2^2 + \dots) f_2^{12} \quad (9.32)$$

We tried to obtain the equivalent formula for $n = 3$ but the calculus was too hard.

Références

- [1] Hajizadeh S., Rezaei-Aghdam A., Poisson-Lie Sigma models over low dimensional real Poisson-Lie groups
- [2] Kosmann-Schwarzbach Y., Lie bialgebras, Poisson Lie Groups and Dressing Transformation
- [3] Vysoký J., Hlavatý , Poisson Lie Sigma Models on Drinfeld double